



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I. De Araneis & Scarabæis Philippenſibus. Ex
MSS. R. P. Geo. Joſ. Camelli. Communicavit
Jacobus Petiver, S. R. S.

De Araneis Crustaceis Philippenſibus.

A Raneos *fila* deducere non ab intrinſeco, tanquam excremento, ut ait *Democritus* & *Plinius*; ſed extrinſecus de ſuo corpore, veluti cortice, aſſerit cum *Aristotele Aldrovandus* & *Forerus*.

^aGAZ. Nat. Verum Anno 1695. (a) Araneos Crustaceos nitidiſſimè
Tab. 26. nigro & flavo picturatos, formæ *Tribuli*, vel potius Pa-
fig. 7. guri, in Horto domeſtico in *Acacia* & *Fagara* cæpi. Unde
Democriti, *Plinij*, *Joſephi Blancani* & aliorum ſententia,
Araneos ab intrinſeco *fila* ſua ducere vindicanda videtur.
^bGAZ. Nat. Alios *teſtaceos* (b) ſexangulares dein Anno inveni ſub-
Tab. 26. ſequenti, *flavo* & *nigro*; alios (c) albo & nigro varia-
fig. 6. tos, & ſex aculeis ſpinofos: *Alium* (d) item poſtea
^cGAZ. Nat. *flavo* & *nigro* variegatum, majorem, & formæ *Lunaris*. an
Tab. 26. *Hoitztocatl Niremberg*. ſeu *Araneus ſpinofus*? morſu de-
fig. 8. mentans, ſuperiore & media corporis portione niger,
^dGAZ. Nat. cætera luteus, formæ aculeatæ & ſimilis *Tribulo*. Aranei
Tab. 26. *crustacei* ictui medetur *Tambalquiſay*, *Camaeſa* & *Piligpoc*.
fig. 5. Nidum vel Ovarium textit electè ſericeum, *flavum* vel
etiam *virentem*, in qua ova reponit *candida*.

1. *Araneus Luzon*. crustaceus, cornu lunulato. GAZ. NATUR. Tab. 26. Fig. 7.

2. *Araneus Luzon*. teſtaceus, trilunatus. GAZOPHYL. NAT. Tab. 26. fig. 6.

3. *Araneus Luzon*. Bovinus. GAZ. NATUR. Tab. 26. fig. 5.

4. *Araneus*

4. *Araneus Luzon. testaceus, angustus, trilunatus.* GAZ. NAT. Tab. 26. fig. 8. 4

De *Araneis* alijs *Luzonensibus*.

5. *Araneus Banayau* 1. *Kamel. Aran. Luzon. mortiferus Smaragdino-aureus.* GAZ. NAT. Tab. 29. fig. 4. Hic ictu soporem & mortem causat Hominibus & Brutis : gaudet *Syringæ* seu *Sambach Arabum* convolutis in folijs habitare : hujus ictui medetur decoctum *Dauvæ* & Ignis. *an Araneus aureus, Lavalava* Nirembergij, stuporem & insaniam inducens ? 5

6. *Araneus parvus brevipes, coloris chermesini, mortiferus.* *Banayau* 2 *Kamel.* 6

7. *Araneus gibbosus Luz. subfuscus, gibbo oblongo, per totum dorsum protracto, elevato & flavo* *Kamel.* GAZ. NATUR. Tab. 29. fig. 12. *Ova* recondit in orbicularibus, flavis & sericeis nidis, *candidula.* 7

8. *Araneus gibbosus alter.* Corpore est *Piso* suppari, dorso proprie in altum gibbum elevato, de reliquo picturatus est aureo fulgente, argenteo relucente & nigro aterrimo. 8

9. *Araneus longipes* 1. *Kamel. Ar. Luz. longipes domesticus.* GAZ NAT. Tab. 50. fig. 1. Corpore est Muscæ tenuiori, domesticus, retiarius, rete varium, quod præda adveniente totum concutit, Communiter Mas & Fæmina in eodem degunt. *Fæmina* habitior est, & una nocte *Ova* excludit, quæ mane in *Piso* pari racemulo congesta ore pendentia gerebat ; die 3 à capturâ pilosa aparuere, 4 magis pilosa, 5 frequentes aranei per vitrum dispersi sunt, remanente inani & candido ovario ab ore matris pendente. 9

10. *Araneus longipes alter* *Kamel. Araneus Luzon. longipes, sem. Ricino æmulans.* GAZ. NATUR. Tab. 50. fig. 4. Habitior est priore, ventre *Piso* pari splendido, & ut semen Ricini picturato, mucronato. 10

11. *Araneus longipes Luzon. 3. Domesticus est, hirsutus & innocuus.* 11

12. *Araneus*

- 12 12. *Araneus longipes Trojanus* five 4. *Bagua & Gam-bagamba Indis*. *Tarantula Luzon.* innocuus. GAZ. NAT. Tab. 50. fig. 2. & A. Corporis mole *Avellanam* æquat, sericeum, albicans, folliculosum, globoso-subplanum ovarium ad ventrem gerens, ex quo mille prodeunt *Araneorum*. Cæterum fuscus, domesticus, innocuus, vespens *Proscara-bæis*. an *Araneus* 10 *Hernandi*?
- 13 13. *Araneus longipes* 5. *Gagamba Luzonenfibus*. Corpore minor, quod cinereo & nigro picturatum est.
- 14 14. *Araneus longipes* 6. *Araneus Luzon.* telarius, argenteo, flavo, & nigro transversim striatus. GAZ. NAT. Tab. 50. fig. 3. Subdialis est. Ejus sericeum album & planum ovarium, in quo ut puto 2000 ovorum alborum latebant. *vid.* Tab. 50. fig. B.
- 15 15. *Araneus Luzon.* oblongus, lutescens, longipes, ventre tenui, geniculo in caudam distincto. GAZ. NATUR, Tab. 43. fig. 14.
- 16 16. *Araneus maximus*, fuscus, pilosus; *Damang Luzon.* Corpore crassitudinis est pollicaris. Degit locis humidis: perniciosus ut *Vipera*. an *Species Tarantulæ*?
- 17 17. *Araneus Luzon.* ex argenteo, flavo & fusco pulchre mixto, pedibus nigris. GAZ. Nat. 29. fig. 6. *Araneus* pulchrè picturatus: Caput, Pectus, corporis binæ striæ laterales, & illa per medium excurrentes, argenteo relucente colore obducta: Binæ striæ inter argenteas excurrentes fuscae, 4 flavis transversim secantibus strijs cancellatæ: pedes longi, nigri.
- 18 18. *Araneus Luzon.* torosus, pilosus, totus fuscus. GAZ. NAT. Tab. 29. fig. 10.
- 19 19. *Araneus Luzon.* torosus, pilosus minor, dorso nigro variegatus.
- 20 20. *Araneus rarus*, niger, splendens; forficulas habens ipso ventre compares; nec primo intuitu facile dictu, quisnam venter, aut quæ sint forficulæ. GAZ. NATUR. Tab. 54. fig. 3.

21. *Araneus venator* Muscarum, in parietum rimis
degens, albo & griseo variegatus: Saltu Muscis insidians. 21
Hispanis Cayman de las Moscas. GAZ. Nat. Tab. 54. 4.
22. *Cotocoto*. *Araneus* est Aquaticus. 22
23. *Laura* vel *Lavalava*. *Araneus* est Domesticus. 23
- De *Scarabæis* Luzonensibus.
- Cantharis* Luzon. *Cetonia* prasino-aurea, luteo-albicantibus maculis variata, ad ventrem ejusdem coloris, sed striatim picturata. GAZOPHYL. NAT. Tab. 26. fig. 9. I GAZ. NAT. Tab. 26. f. 9.
- Cantharis* Luzon. pulcherrima, vaginâ integrâ non fissâ, quinis aureo-cyaneis maculis ornatâ, ipsa Havo-aurea; venter medius superne smaragdino-aureus & tribus aureo-cyaneis maculis notatus ut vagina dorsi: Inferior venter aureis & auréo-virentibus strjis transversim virgatus est. 2 G. N. 37. 7.
- Stockerus* Luzon. viridis, scapulâ maculisq; purpureis. GAZ. NATUR. Tab. 37. fig. 7.
- Cantharis* Luzon. *minima*, magnitudinis est hujatis Cicindelæ, smaragdino-aurea, capite & pedibus cyaneis, collo luteo, corniculis plumatis. *Indis* Alibamban & Parapan. 3
- Cantharis* Catunduana & Bicolana *major*, palmo longior, auriluca, smaragdina. *Indis* Olaniban & Basingoban. 4
- Cantharis* oblonga communis, nitidissime tota, aureo-viridis, *Camarinensium*. Diau, Giao, Laniban & Salibat *Indis*. GAZ. NAT. Tab. 48. fig. 5. 5 G. N. 48. 5.
- Cantharis* cæruleo-aurea, elytris ad medium luteo-aureis maculis notatis. GAZ. NAT. Tab. 49. 7. 7 G. N. 49. 2.
- Cantharis* merè viridis, *Indis* Ticpaolong. GAZ. NAT. Tab. 41. fig. 2.
- Cantharis* auriluca, viridante & rubente striatim varia. *Cochinella* Luzon. *major*, è cupreo viridique eleganter striata. GAZ. NAT. Tab. 41. fig. 3. 8 G. N. 41. 3.
- Cantharis* Luzonis nona & maxima, perpolita, tota multò splendidior quintâ, colore aureo-smaragdino per-G. N. 28. 3. 9
- Z z
- petuante

petuante. Differt a Quintâ, capite, quod longius; mustaciis, qui curti & subplani; scapulari, quod longius angulose protensum, & pedibus, quorum ultima articulatio spinosa, uropygio, quod coloris est cuprei. *Oculi nigri sunt.*

Scarabæus pectinatus Luzon. aureo-smaragdinus. GAZ. NAT. Tab. 28. fig. 3.

9 *Capricornus Luzonicus, seu Carabus, five Cerambix, facie G. N. 37. 6. Tauri. Antennis articulatim nodosis; coloris in fusco, cinereo-lutescente undulatus, capite valido, facie Tauri, duobus majoribus & aduncis dentibus instructus. GAZ. NAT. Tab. 37. fig. 6.*

10 *Capricornus alter, atratus est & striatus. GAZ. NAT. G. N. 37. 12. Tab. 37. fig. 12.*

11 *Capricornus Luzon. 3. Totus ex atro & pomi aurei colore picturatus est. GAZ. NAT. Tab. 28. fig. 5.*

12 *Capricornus seu Cerambix Luzon. maxima. GAZ. NAT. G. N. 47. 1. Tab. 47. fig. 1.*

13 *Buceros Nasicornis 1. Luzon. Hoang. GAZ. NAT. Tab. G. N. 49. 8. 49. fig. 8. & Kamel. in Append. Ray Hist. Plant. V. 3. p. 44.*

14 *Buceros Nasicornis alter Bogang Luzon. GAZ. NAT. G. N. 29. 2. Tab. 29. fig. 2 & Kamel in App. Ray. p. 44.*

15 *Melolonthes seu Cantharis merè viridis major. Indis Salaguinto. Germanis Gruner, May-Koffer, & S. Joannis Koffer.*

16 *Melolonthes seu Cantharis merè viridis minor. Indis G. N. 41. 2. Ticipaolong, Germ. Gold Koffer, id No. 7. G. N. 41. 2.*

N. B. *Inveniuntur & Flavæ, Rubæ, Ceruleæ, & variè picturata.*

17 *Scarabæus arboreus, seu Bruchus Majalis sublustis, in G. N. 38. 11. fusco cinereis punctis notatus, Europæo compar & major. Indis Salibamban & Salagubat, Germ. Mayen-Koffer & Baum-Koffer. GAZ. NAT. Tab. 38. fig. 11.*

18 *Scarabæus arboreus 2. luteo-fuscus, tenuissimè striatus. G. N. 38. 6. GAZ. NAT. Tab. 38. 6.*

19 *Scarabæus arboreus 3 subplanus, sublustis, luride èrugi- G. N. 38. 5. nose maculosus. G. N. Tab. 38. fig. 5.*

Scarabæus atratus, elytris striatis, forcipulâ validâ bifurcatâ, quâ arbores atterebat, donatus. G. N. Tab. 42. 3. 20

Scarabæus mirabilis, caput lato & denticulato scuto contegente. Gaudens *Scilla*. GAZ. NAT. Tab. 42. fig. 2. 21
G. N. 42. 2

Scarabæus pilularis torosus, niger, *Luzonis* minimus. 22
Germ. Ross-Koffer. GAZ. NAT. Tab. 42. fig. 5. G. N. 42. 5

Scarabæus Luz. elasticus, vaginis striatis. GAZ. NAT. 23
Tab. 27. fig. 10. G. N. 27. 10.

Scarabæus saltator Luz. *Germ.* Springer. *Indis* Dangdo. Hujas *Europæo* major est, sublustris, coloris terreo-fusci, non ater ut ille; ad Pectus habet solidum Stimulum, quo supinè positus nititur in altum vibrari. Elytra striata, caput parvulum.

Scarabæus Saltator, seu ille parvus, oblongus, niger, qui supinè positus e latere intenso in altum vibratur; in *Luzone* sordidè niger est & sesquiuncialis. *Hisp.* Batecavezzi. 24
Germ. Springer & Schuester. *Indis* Baldoc bondoc. *Tagal.* Dangdo.

Scarabæus Saltator alter parvulus. *Europæo* minor est, luteus, & ad extremum niger. 25

Proscarabæus nocturnus domesticus *Kameli*.

Blatta domestica Luzon. nocturna. GAZ. NAT. Tab. 48. fig. 9. 26
Cucca & *Cuccaracha* *Hispan.* *Schioabem* *Germ.* *Lusit.* Barata. *Ipis* & *Bangcocang Indis.* hujus est Species *Gukiana*. G. N. 48. 9.

Insectum est malè olens, elytris tenuibus, omnia depascens & arrodens. *Mas* coloris grisei, aliquibus nigris punctis variegatus. *Femina* tota coloris castanei, & minor mare, ova excludens oblongo-compressa ad latus acuta. *Mas* & *Femina* depositâ senectâ candidi sunt & tenelli. Utuntur *Indi* cinere poto in Astmate, confusis cum Saccharo, & impositis in Cancro & Panaritio & ulceribus suppurandis. Assos No. 6 vel 8. exhibent *Pueræ* lumbricis laborantibus, quos enecant.